

MEETING ABSTRACT

Open Access

Respiratory mechanics influence VO_2 max in acute hypoxia in females

Teruhiro Hanamaru¹, Tsuyoshi Yoshikawa¹, Takeshi Nishiyasu², Takeshi Ogawa^{1*}

From 15th International Conference on Environmental Ergonomics (ICEE XV)
Portsmouth, UK. 28 June - 3 July 2015

Introduction

The magnitude of decrease in maximal oxygen uptake (VO_2 max) in hypoxia (H) is more pronounced in male endurance athletes. In these subjects, high pulmonary ventilation (V_E) could be beneficial in maintaining VO_2 max in H [1]. Because females have smaller chest walls and narrower airways than males, V_E during intensive exercise is mechanically limited [2]. Thus, it is hypothesised that in females, respiratory response and mechanics influenced the magnitude of decrease in VO_2 max in H relative to males, despite lower VO_2 max in females than in males. To test this hypothesis, we studied 22 healthy males and females as they performed an exhaustive cycling test in H and normoxia (N) conditions.

Methods

Twenty-two healthy males ($n = 12$; VO_2 max: 51 (7) $\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$, age; 21 (2) yr, stature; 172 (3) cm, mass; 66 (6) kg) and females ($n = 10$; VO_2 max = 44 (6) $\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$, age; 21 (1) yr, stature; 159 (4) cm, mass; 53 (8) kg) performed the incremental cycle exercise test until exhaustion under N (20.9 % O_2) and H (15.0 % O_2) conditions. During the exercise test, we measured VO_2 max and V_E max. To mechanically assess the respiratory work, we measured transpulmonary pressure by subtracting mouth pressure from esophageal pressure and calculated work of breathing (WOB) as the integrated area of the Ptp-volume loop.

Results

The percentage decrease in VO_2 max in H (% $d\text{VO}_2$ max) tended to be larger in females than in males (-16% in males and -21% in females, $p < 0.06$). V_E/VO_2

was significantly ($p < 0.05$) higher in females than males, and it was significantly ($p < 0.01$) higher in H than in N in both genders. In females only, the % $d\text{VO}_2$ max in H was significantly correlated to the extent of change in V_E max ($r = 0.79$, $p < 0.05$). In comparison with N, WOB/V_E in H tended to be lower in males (-13.1%) whereas it was 14.6% higher in females (not significant). Furthermore, in females, the % $d\text{VO}_2$ max in H was significantly correlated to WOB/V_E in H ($r = -0.76$, $p < 0.05$).

Discussion

These results suggest that females have lower ventilatory mechanical efficiency than males, and V_E is one of the factors causing this decrease in VO_2 max in H. Further, there is a possibility that the oxygen demand at the respiratory muscles greatly increases against the increase in V_E in H. Thus, high respiratory muscle work compromises blood flow to the active muscles [3], thereby limiting their peak work rate and VO_2 max in H.

Conclusion

Our findings demonstrated that in females, the respiratory muscle work efficiency affected the decrease in VO_2 max in H, a decrease which tended to be larger in females than in males, despite the lower VO_2 max in females compared with males.

Authors' details

¹Department of Physical and Health Education, Osaka Kyoiku University, Japan. ²Institute of Health and Sports Science, University of Tsukuba, Japan.

Published: 14 September 2015

References

1. Ogawa T, Calbet JA, Honda Y, Fujii N, Nishiyasu T: The effects of breathing a helium-oxygen gas mixture on maximal pulmonary ventilation and maximal oxygen consumption during exercise in acute moderate hypobaric hypoxia. *Eur J Appl Physiol* 2010, **110**(4):853-861.

* Correspondence: ogawat@cc.osaka-kyoiku.ac.jp

¹Department of Physical and Health Education, Osaka Kyoiku University, Japan

Full list of author information is available at the end of the article

2. McClaran SR, Harms CA, Pegelow DF, Dempsey JA: **Smaller lungs in women affect exercise hyperpnea.** *J Appl Physiol* 1998, **84**(6):1872-1881.
3. Harms CA, Babcock MA, McClaran SR, Pegelow DF, Nickele GA, Nelson WB, *et al*: **Respiratory muscle work compromises leg blood flow during maximal exercise.** *J Appl Physiol* 1997, **82**(5):1573-1583.

doi:10.1186/2046-7648-4-S1-A64

Cite this article as: Hanamaru *et al.*: Respiratory mechanics influence VO_2 max in acute hypoxia in females. *Extreme Physiology & Medicine* 2015 **4**(Suppl 1):A64.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

